

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

CLiKZiN 12.5 mg/ml Pour-On Suspension for Sheep

### **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each ml contains:

**Active substance:**

Dicyclanil 12.5 mg

**Excipients:**

Quinoline yellow (E104)	0.05 mg
Patent blue V (E131)	0.05 mg
Methyl parahydroxybenzoate (E218)	1.50 mg
Propyl parahydroxybenzoate	3.00 mg
Butylated hydroxytoluene (E321)	0.50 mg

For the full list of excipients, see section 6.1

### **3. PHARMACEUTICAL FORM**

Pour-on suspension.  
Green coloured.

### **4. CLINICAL PARTICULARS**

#### **4.1 Target species**

Sheep

#### **4.2 Indications for use, specifying the target species**

Prevention of blowfly strike on sheep due to *Lucilia sericata*.

#### **4.3 Contraindications**

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

#### **4.4 Special warnings for each target species**

The product is best applied before an anticipated blowfly challenge, or when a blowfly challenge is identified on or in the vicinity of the farm. Established strikes may require a separate treatment with a knockdown insecticide.

It is recommended that animals with dirty back-ends are dagged or crutched prior to application. If dagging or crutching is undertaken in the weeks following application, these animals should be re-treated, otherwise protection could be lost.

Do not apply during heavy rainfall, or when such conditions are expected. The resulting wash out may reduce the protection period.

#### 4.5 Special precautions for use

i). Special precautions for use in animals

None.

ii). Special precautions to be taken by the person administering the veterinary medicinal product to animals

Redness and irritation may develop after skin or eye contact with the product.

- Contact with skin and eyes should be avoided.
- Personal protective equipment consisting of synthetic rubber gloves and PVC trousers should be worn when handling the product.
- In case of skin contact remove contaminated clothing and thoroughly wash the affected parts of the body with soap and water.
- In case of eye contact, wash immediately with clean water.
- Always wash hands and exposed skin with soap and water after work.
- Do not eat, drink or smoke whilst using the product.
- It is good agricultural practice to minimise handling of sheep after treatment. If you need to handle sheep within 2 months after treatment, wear synthetic rubber gloves and long trousers or coveralls. If sheep are wet wear waterproof trousers.

#### Special precautions for the protection of the environment

Treated sheep **must** be kept away from watercourses for at least one hour after treatment. There is a **serious** risk to aquatic life if this advice is not followed.

iii). Other precautions

#### 4.6 Adverse reactions (frequency and seriousness)

Target Species': Sheep

None known.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a

veterinarian, to either the marketing authorisation holder or the national competent authority via the national reporting system. See also the last section of the package leaflet for contact details.

#### **4.7 Use during pregnancy, lactation or lay**

The safety of the veterinary medicinal product has not been established during pregnancy or lactation.

Use only according to the benefit/risk assessment by the responsible veterinarian.

Laboratory studies in rats and rabbits have not produced any evidence of teratogenic, foetotoxic or maternotoxic effects.

#### **4.8 Interaction with other medicinal products and other forms of interaction**

None known.

#### **4.9 Amount(s) to be administered and administration route**

For external use only.

The veterinary medicinal product is applied according to the following recommendations:

<b>Bodyweight (kg)</b>	<b>Dose Volume (ml)</b>
<b>10-20</b>	<b>20</b>
<b>21-30</b>	<b>24</b>
<b>31-50</b>	<b>30</b>
<b>&gt;50</b>	<b>36</b>


(Guide dose volumes correspond to 0.7–2 ml [7.5–25 mg dicyclanil] per kg bodyweight.)

To ensure administration of a correct dose, body weight should be determined as accurately as possible; accuracy of the dosing device should be checked.

If animals are to be treated collectively rather than individually, they should be grouped according to their bodyweight and dosed accordingly, in order to avoid under- or overdosing.

Recommended for treatment in sheep 3 weeks post shearing. Shake the container well before use.

The veterinary medicinal product must be applied with a manual or automatic dosing gun, fitted with a spray nozzle, which guarantees the correct spreading of the product on the fleece. Best results will be achieved by holding the gun approximately 45 cm from the sheep during application. Apply as a fan spray using a 4 stroke method as shown in the pictogram along the spine of the animal in bands at least 10 cm wide from the middle of the shoulders and in an arc around the crutch and tail. Half the dose should be applied along the spine with the remainder over the tail and crutch area.



The veterinary medicinal product should be administered before or at the start of predicted fly activity but is also suitable for use during the fly season.

The veterinary medicinal product will protect against fly strike for 8 weeks. It is good agricultural practice to check animals regularly for fly strike.

#### **4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary**

An overdose of up to at least 20 times the recommended dose does not lead to any signs of local or systemic intolerance.

No antidote is known.

#### **4.11 Withdrawal period(s)**

Meat and offal: 7 days

Not authorised for use in animals producing milk for human consumption.

## **5. PHARMACOLOGICAL PROPERTIES**

**Pharmacotherapeutic group:** Other ectoparasiticides for topical use

**ATC Vet Code:** QP53AX24.

### **5.1 Pharmacodynamic properties**

Dicyclanil prevents the moult from the first to the second larval instar of *Lucilia* spp. It is less effective against later larval stages and does not have any adulticide action. The mode of action of dicyclanil is believed to be similar to that of the triazine compounds.

### **5.2 Pharmacokinetic particulars**

In studies with a more concentrated 5% (w/v) formulation, after 7 days post dosing, approximately 5% of the dose was absorbed and eliminated in urine and faeces. Peak blood levels were observed between 12 and 48h post dose, accounting for <0.025 mg dicyclanil equivalents/kg.

In experimental metabolism studies, absorbed radioactivity was widely distributed throughout the body. Longest half lives were found in liver and kidney being 13 and 10 days respectively. In muscle, fat and wool, unchanged dicyclanil was found to be the major residue, whereas in liver and kidney the descyclopropyl dicyclanil was found to be the major residue together with unchanged dicyclanil.

In field residue depletion studies on sheep with the veterinary medicinal product, residue levels at day 7 were very low (at maximum 31.9 and 30.7 µg/kg in liver and kidney, respectively, and no quantifiable residues found in muscle or fat), indicating minimal systemic absorption.

### **5.3 Environmental properties**

The use of the veterinary medicinal product has harmful effects on dung flies and beetles.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Quinoline yellow (E 104)  
Patent blue V (E 131)  
Methyl parahydroxybenzoate (E 218)  
Propyl parahydroxybenzoate  
Butylated hydroxytoluene (E 321)  
Polysorbate 20  
Acrylic acid copolymer  
Disodium edetate  
Distilled monoglycerides

Triglycerides, medium-chain  
Propylene glycol  
Sodium hydroxide  
Purified water

## **6.2 Major Incompatibilities**

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

## **6.3 Shelf life**

Shelf life of the veterinary medicinal product as packaged for sale: 3 years  
Shelf life after first opening the immediate packaging: 1 year

## **6.4 Special precautions for storage**

Protect from frost.  
Store in the original container.  
Keep the container tightly closed, away from food, drink and animal feedstuffs.  
Protect from direct sunlight.

## **6.5 Nature and composition of immediate packaging**

The pack is composed of pigmented white opaque polyethylene back pack container with blue polypropylene screw cap, containing 0.8, 2.2 or 5 litres of finished product.

Not all pack sizes may be marketed.

## **6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products**

Medicines should not be disposed of via wastewater.

The product should not enter water courses as dicyclanil may be dangerous for fish and other aquatic organisms.

Do not contaminate ponds or other waterways with product or empty containers.

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

## **7. MARKETING AUTHORISATION HOLDER**

Elanco GmbH  
Heinz-Lohmann Strasse 4  
Grodan  
27472 Cuxhaven  
Germany

**8. MARKETING AUTHORISATION NUMBER**

Vm 52127/5094

**9. DATE OF FIRST AUTHORISATION**

15 September 2010

**10. DATE OF REVISION OF THE TEXT**

May 2025

**11. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS**

Veterinary medicinal product subject to prescription.

*Gavin Hall*

Approved: 22 May 2025