

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Nobivac Myxo-RHD PLUS lyophilisate and solvent for suspension for injection for rabbits

### **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each dose (0.2 ml or 0.5 ml) of reconstituted vaccine contains:

#### **Active substances:**

Live myxoma vectored RHD virus strain 009:  $10^{3.0} - 10^{5.8}$  FFU\*

Live myxoma vectored RHD virus strain MK1899:  $10^{3.0} - 10^{5.8}$  FFU\*

\*Focus Forming Units

For the full list of excipients, see section 6.1.

### **3. PHARMACEUTICAL FORM**

Lyophilisate and solvent for suspension for injection.

Lyophilisate: off-white or cream-coloured pellet.

Solvent: clear colourless solution.

### **4. CLINICAL PARTICULARS**

#### **4.1 Target species**

Rabbits.

#### **4.2 Indications for use, specifying the target species**

For active immunisation of rabbits from 5 weeks of age onwards to reduce mortality and clinical signs of myxomatosis and rabbit haemorrhagic disease (RHD) caused by classical RHD virus (RHDV1) and RHD type 2 virus (RHDV2).

Onset of immunity: 3 weeks.

Duration of immunity: 1 year.

#### **4.3 Contraindications**

None.

#### 4.4 Special warnings for each target species

Vaccinate healthy animals only.

High levels of maternally derived antibodies against myxoma virus and/or RHD virus can potentially reduce the efficacy of the product. To ensure the full duration of immunity, vaccination from 7 weeks of age is advised in this case.

Rabbits that have been vaccinated previously with another myxomatosis vaccine, or that have experienced natural myxomatosis infection in the field, may not develop an adequate immune response against rabbit haemorrhagic disease following vaccination.

#### 4.5 Special precautions for use

Special precautions for use in animals:

Not applicable.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Not applicable.

Special precautions for the protection of the environment:

Not applicable.

Other precautions:

Not applicable.

#### 4.6 Adverse reactions (frequency and seriousness)

Rabbits:

Common (1 to 10 animals / 100 animals treated):	Hyperthermia <sup>1</sup> , Injection site swelling. <sup>2</sup>
Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Injection site necrosis <sup>3</sup> , injection site scab <sup>3</sup> , injection site crust <sup>3</sup> , injection site hair loss <sup>3</sup> . Hypersensitivity reaction <sup>4</sup> . Myxomatosis <sup>5</sup> . Anorexia, lethargy.

<sup>1</sup> Transient temperature increase of 1 – 2 °C

<sup>2</sup> A small, non-painful swelling (maximum 2 cm diameter) within the first two weeks after vaccination. The swelling will resolve completely by 3 weeks after vaccination.

<sup>3</sup> In pet rabbits.

<sup>4</sup> Sometimes fatal.

<sup>5</sup> Mild clinical signs of myxomatosis may occur within 3 weeks of vaccination.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a veterinarian, to either the marketing authorisation holder or the national competent authority via the national reporting system. See the package leaflet for respective contact details.

#### **4.7 Use during pregnancy, lactation or lay**

Pregnancy:

Can be used during pregnancy.

Fertility:

No safety study on the reproductive performance has been conducted in male rabbits (bucks). Therefore, the vaccination of breeding bucks is not recommended.

#### **4.8 Interaction with other medicinal products and other forms of interaction**

No information is available on the safety and efficacy of this vaccine when used with any other veterinary medicinal product. A decision to use this vaccine before or after any other veterinary medicinal product therefore needs to be made on a case by case basis.

#### **4.9 Amount(s) to be administered and administration route**

Subcutaneous use.

Primary vaccination:

Administer one dose to rabbits from 5 weeks of age onwards.

Revaccination:

Revaccinate annually.

Ensure that the lyophilisate is completely reconstituted before use.

Reconstituted product: off-pink or pink coloured suspension.

Single dose vial

Reconstitute a single dose vial containing lyophilisate with 0.5 ml of the supplied solvent. Administer the total contents of the vial.

Multi-dose vial (50 doses)

Reconstitute a multi-dose vial containing lyophilisate with 10 ml of the supplied solvent. Administer 0.2 ml per animal.

For proper reconstitution of the multi-dose vial, use the following procedure:

1. Add 1 – 2 ml of solvent to the 50-dose vaccine vial and ensure that the lyophilisate is fully dissolved.
2. Withdraw the reconstituted vaccine concentrate from the vial and inject it back into the solvent vial.
3. Ensure that the resulting vaccine suspension in the solvent vial is properly mixed.
4. Use the vaccine suspension within 4 hours of reconstitution. Any reconstituted vaccine remaining at the end of this time should be discarded.

#### **4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary**

In addition to the adverse reactions observed after single dose vaccination, a mild swelling of the local lymph nodes may be observed within the first 3 days after administration of a ten-fold overdose.

#### **4.11 Withdrawal period(s)**

Zero days.

### **5. IMMUNOLOGICAL PROPERTIES**

**Pharmacotherapeutic group:** Immunologicals for *Leporidae*, live viral vaccine for rabbits

**ATCvet code:** QI08AD02.

The vaccine is intended to stimulate immunity against myxoma virus and rabbit haemorrhagic disease viruses in rabbits.

The vaccine strains are myxoma viruses expressing the capsid protein gene of classical or type 2 RHD viruses. As a consequence, rabbits are immunised against myxoma virus and both classical and type 2 RHD viruses.

After infection with virulent field myxoma virus some vaccinated animals may develop a few very small swellings, especially on hairless places of the body, which quickly form scabs. These scabs usually disappear within 2 weeks. The scabs are only observed in animals with active immunity and have no influence on the general health, appetite or behaviour of the rabbit. Recent or latent infection with field myxoma virus seems to play a role in the development of the mild clinical signs of myxomatosis that may occur within 3 weeks after vaccination.

### **6. PHARMACEUTICAL PARTICULARS**

#### **6.1 List of excipients**

Lyophilisate:

Hydrolysed gelatine  
Pancreatic digest of casein  
Sorbitol  
Disodium phosphate dihydrate

Solvent:

Disodium phosphate dihydrate  
Potassium dihydrogen phosphate  
Water for injections

## 6.2 Major incompatibilities

Do not mix with any other veterinary medicinal product, except solvent supplied for use with the veterinary medicinal product.

## 6.3 Shelf life

Shelf life of the veterinary medicinal product (lyophilisate) as packaged for sale: 2 years.

Shelf life of the solvent as packaged for sale: 4 years.

Shelf life after reconstitution according to directions: 4 hours.

## 6.4 Special precautions for storage

### Lyophilisate:

Store in a refrigerator (2 °C – 8 °C).

Do not freeze.

Protect from light.

### Solvent:

No special precautions for storage.

## 6.5 Nature and composition of immediate packaging

### Lyophilisate:

Type I clear glass vial of 1 or 50 doses closed with a chlorobutyl rubber stopper and aluminium cap.

### Solvent:

Type I clear glass vial of 0.5 ml or 10 ml closed with a bromobutyl rubber stopper and aluminium cap.

### Pack sizes:

- Plastic box with 5 x 1 dose vial of vaccine and 5 vials containing 0.5 ml of solvent.
- Plastic box with 25 x 1 dose vial of vaccine and 25 vials containing 0.5 ml of solvent.
- Cardboard box with 10 x 50 dose vials of vaccine; and cardboard box with 10 x 10 ml vials of solvent.

Not all pack sizes may be marketed.

## 6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products

Medicines should not be disposed of via wastewater.

Any unused veterinary medicinal product or waste materials derived from such medicinal product should be disposed of in accordance with local requirements.

## **7. MARKETING AUTHORISATION HOLDER**

MSD Animal Health UK Limited  
Walton Manor  
Walton  
Milton Keynes  
MK7 7AJ

## **8. MARKETING AUTHORISATION NUMBER**

Vm 01708/5050

## **9. DATE OF FIRST AUTHORISATION**

05 December 2019

## **10. DATE OF REVISION OF THE TEXT**

May 2024

## **PROHIBITION OF SALE, SUPPLY AND/OR USE**


Any person intending to manufacture, import, possess, distribute, sell, supply and use this veterinary medicinal product must first consult the relevant competent authority on the current vaccination policies, as these activities may be prohibited in a country on the whole or part of its territory pursuant to national legislation.

## **11. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS**

Veterinary medicinal product subject to prescription.

Find more product information by searching for the 'Product Information Database' or 'PID' on [www.gov.uk](http://www.gov.uk).

Approved 14 May 2024



J. Hunter.