

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

TUR-3

### **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

#### Active substances:

Inactivated Paramyxovirus Type 3, at least .....40 HAIU

Inactivated Newcastle Disease virus, at least .....50 PD<sub>50</sub>

Inactivated Turkey Rhinotracheitis virus, at least..... 9 EU

#### Adjuvant:

Paraffin oil.....170 to 186 mg

#### Excipient:

Thiomersal..... 15 µg

1 HAIU: q.s. to obtain a mean haemagglutination inhibiting antibody titre of 1 in vaccinated animals

1 EU: q.s. to obtain a positive serum by ELISA in one vaccinated animal

50 PD<sub>50</sub> (50% protective dose) is performed in chickens.

For full list of excipients, see section 6.1

### **3. PHARMACEUTICAL FORM**

Suspension for injection.

### **4. CLINICAL PARTICULARS**

#### **4.1 Target species**

Future breeder turkeys.

#### **4.2 Indications for use, specifying the target species**

For active immunisation of future breeder turkeys:

- as booster vaccination after priming with live vaccines against Newcastle Disease and Turkey Rhinotracheitis to reduce mortality and clinical signs of Newcastle Disease and to induce a specific seroconversion against Newcastle Disease and Turkey Rhinotracheitis in vaccinated birds throughout the laying period

- as vaccination against Paramyxovirus Type 3 to reduce the decrease in egg production, as demonstrated by challenge at peak lay, and to induce a specific seroconversion against paramyxovirus type 3 throughout the laying period.

Onset of immunity: 4 weeks after the first injection.

Duration of immunity: one laying period (demonstrated by serology).

#### **4.3 Contra-indications**

None.

#### **4.4 Special warnings for each target species**

None.

#### **4.5 Special precautions for use**

##### **(i) Special precautions for use in animals**

Vaccinate only healthy animals.

Apply usual aseptic procedures.

Do not use syringes with natural rubber or butyl elastomer pistons.

Do not keep partly used containers - use immediately after opening. Remove from the refrigerator and allow to reach room temperature before using.

Shake well before use.

TUR-3 may be used with automatic vaccinating equipment. Equipment including needles and syringes must be sterile before use.

##### **(ii) Special precautions to be taken by the person administering the veterinary medicinal product to the animals**

###### To the user:

This product contains mineral oil. Accidental injection/self injection may result in severe pain and swelling, particularly if injected into a joint or finger, and in rare cases could result in the loss of the affected finger if prompt medical attention is not given.

If you are accidentally injected with this product, seek prompt medical advice even if only a very small amount is injected and take the package leaflet with you.

If pain persists for more than 12 hours after medical examination, seek medical advice again.

###### To the physician:

This product contains mineral oil. Even if small amounts have been injected, accidental injection with this product can cause intense swelling, which may, for example, result in ischaemic necrosis and even loss of a digit. Expert, PROMPT, surgical attention is required and may necessitate early incision and irrigation of the injected area, especially where there is involvement of finger pulp or tendon.

#### **4.6 Adverse reactions (frequency and seriousness)**

None.

#### **4.7 Use during pregnancy, lactation or lay**

This product is not recommended for administration to birds 'in lay'.

#### **4.8 Interaction with other medicinal products and other forms of interaction**

No information is available on the safety and efficacy of this vaccine when used with any other veterinary medicinal product. A decision to use this vaccine before or after any other veterinary medicinal product therefore needs to be decided on a case by case basis.

#### **4.9 Amounts to be administered and administration route**

##### **Posology:**

One 0.3-ml dose.

##### **Method and route of administration:**

Shake well before use.

Intramuscular route.

Primary vaccination: one injection 8 to 10 weeks before the beginning of lay.

Booster vaccination: one injection 2 to 4 weeks before the beginning of lay.

#### **4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary**

In the double dose overdose study no abnormal local or systemic reactions occurred.

#### **4.11 Withdrawal periods**

Zero hours/days.

### **5. IMMUNOLOGICAL PROPERTIES**

The vaccine induces a specific serological response against Newcastle disease, Turkey Rhinotracheitis and Paramyxovirus Type 3, in vaccinated animals that persists for 21 weeks post completion of the vaccination course. Studies have demonstrated that the vaccine can be used as an aid in the prevention of 'egg drop' caused by Paramyxovirus Type 3 challenge.

##### **ATC Vet Code:**

QI01CA02

### **6. PHARMACEUTICAL PARTICULARS**

#### **6.1 List of excipients**

Thiomersal

Paraffin oil

Ester of fatty acids and polyols

## **6.2 Major incompatibilities**

Do not mix with any other veterinary medicinal product.

## **6.3 Shelf-life**

Shelf-life of the veterinary medicinal product as packaged for sale: 24 months.

## **6.4 Special precautions for storage**

Store between +2°C and +8 °C, protected from light. Do not freeze.

Once broached, use immediately.

## **6.5 Nature and composition of immediate packaging**

500-dose (polypropylene) bottle.

500-dose (polypropylene) bottle, 10 bottle package.

1,000-dose (polypropylene) bottle.

1,000-dose (polypropylene) bottle, 10 bottle package.

## **6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products, if appropriate**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

## **7. MARKETING AUTHORISATION HOLDER**

Boehringer Ingelheim Vetmedica GmbH  
Binger Strasse 173  
55218 Ingelheim am Rhein  
Germany

## **8. MARKETING AUTHORISATION NUMBER**

Vm 61700/5024

## **9. DATE OF FIRST AUTHORISATION**

23 May 1997

## **10. DATE OF REVISION OF THE TEXT**

December 2025

## **ADDITIONAL INFORMATION**

This is a Limited Marketing Authorisation and the limitations of the product are reflected in the claims.

All suspected adverse reactions and any suspected lack of efficacy should be reported to 01344 746957 at Boehringer Ingelheim Animal Health UK Ltd., Ellesfield Avenue, Bracknell, Berkshire, RG12 8YS, UK.

Further information on this product and its supporting data can be found on **[HTTP://WWW.VMD.GOV.UK/PRODUCTINFORMATIONDATABASE](http://www.vmd.gov.uk/productinformationdatabase)**.

*Gavin Hall*  
Approved: 02 December 2025