SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Panomec Injection for Cattle, Sheep and Pigs

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each ml contains:

Active substance:

Ivermectin 10 mg

Excipients:

Qualitative composition of excipients and other constituents		
Glycerol formal		
Propylene glycol		

A clear, pale, straw-coloured liquid.

3. CLINICAL INFORMATION

3.1 Target species

Cattle, sheep and pigs.

3.2 Indications for use for each target species

The veterinary medicinal product is indicated for the effective treatment and control of the following harmful parasites of cattle, sheep and pigs:

Cattle

			Inhibited
PARASITE	Adult	L4	L4
Gastrointestinal roundworms:			
Ostertagia lyrata	•	•	
Ostertagia ostertagi	•	•	•
Cooperia oncophora	•	•	
Cooperia pectinata	•	•	
Cooperia punctata	•	•	
Haemonchus placei	•	•	
Trichostrongylus axei	•	•	
Trichostrongylus colubriformis	•	•	
Bunostomum phlebotomum	•	•	
Oesophagostomum radiatum	•	•	
Strongyloides papillosus	•		
Nematodirus helvetianus	•		
Nematodirus spathiger	•		

Trichuris spp.

Lungworms:

Dictyocaulus viviparus • •

Eye worms: Thelazia spp

Warbles:

Hypoderma bovis

H. lineatum

Mange mites:

Psoroptes ovis

Sarcoptes scabiei var. bovis

Sucking lice:

Linognathus vituli

Haematopinus eurysternus

Solenopotes capillatus

The veterinary medicinal product may also be used as an aid in the control of biting lice (*Damalinia bovis*) and the mange mite (*Chorioptes bovis*), but complete elimination may not occur.

Persistent activity

The veterinary medicinal product given at the recommended dosage of 0.2 mg per kg bodyweight controls re-infection with:

Parasite	No. Of Days After Treatment
Barbers pole worm – <i>Haemonchus placei</i>	14
Small intestinal worm – <i>Cooperia</i> spp.	14
Hairworm – <i>Trichostrongylus axei</i>	14
Brown stomach worm – Ostertagia ostertagi	21
Nodular worm – Oesophagostomum radiatum	21
Lungworm – <i>Dictyocaulus viviparus</i>	28

The timing of treatment should be based on epidemiological factors and should be customised for each individual farm. A dosing program should be established by a qualified professional person.

Sheep

·			Inhibited
PARASITE	Adult	L4	L4
Gastrointestinal roundworms:			
Ostertagia circumcincta	•	•	•
O. trifurcata	•	•	
Haemonchus contortus	•	•	•
Trichostrongylus axei	•		
T. colubriformis	•	•	
T. vitrinus	•		
Cooperia curticei	•	•	
Oesophagostomum columbianum	•	•	

O. venulosum
Nematodirus filicollis
Chabertia ovina
Trichuris ovis
Lungworms
Dictyocaulus filaria
Protostrongylus rufescens
Nasal bots
Oestrus ovis
Mange mites
Psoroptes ovis*

Pigs

PARASITE	Adult	L4
Gastrointestinal roundworms		
Ascaris suum	•	•
Hyostrongylus rubidus	•	•
Oesophagostomum spp	•	•
Strongyloides ransomi *	•	
<u>Lungworms</u>		
Metastrongylus spp	•	
<u>Lice</u>		
Haematopinus suis		
Mange mites		
Sarcoptes scabei var. suis		
* Includes somatic larval stages		

3.3 Contraindications

The veterinary medicinal product is not for intramuscular or intravenous use. Do not use in cases of hypersensitivity to the active substance or any of the excipients.

3.4 Special warnings

In sheep, treatment of psoroptic mange (sheep scab) with one injection is not recommended because, although a clinical improvement may be seen, elimination of all mites may not occur.

Care should be taken to avoid the following practices because they increase the risk of development of resistance and could ultimately result in ineffective therapy:

- too frequent and repeated use of anthelmintics from the same class, over an extended period of time.
- underdosing, which may be due to underestimation of bodyweight,

^{*}For the treatment and control of sheep scab, two injections with a seven-day interval are required to treat clinical signs of scab and to eliminate mites.

misadministration of the product, or lack of calibration of the dosing device.

Suspected clinical cases of resistance to anthelmintics should be further investigated using appropriate tests (e.g. Faecal Egg Count Reduction Test). Where the results of the test(s) strongly suggest resistance to a particular anthelmintic, an anthelmintic belonging to another pharmacological class and having a different mode of action should be used.

Resistance to macrocyclic lactones (which includes ivermectin) has been reported in *Teladorsagia* spp. in sheep and in *Cooperia* spp. in cattle within the EU. Therefore, the use of this product should be based on local (regional, farm) epidemiological information about susceptibility of nematodes and recommendations on how to limit further selection for resistance to anthelmintics.

3.5 Special precautions for use

Special precautions for safe use in the target species:

The veterinary medicinal product has been formulated specifically for use in these target species. It should not be used in other species as severe adverse reactions, including fatalities in dogs, may occur.

Sheep scab (*Psoroptes ovis*) is an extremely contagious external parasite of sheep. Following treatment of infected sheep, great care must be taken to avoid re-infestation, as mites may be viable for up to 15 days off the sheep. It is important to ensure all sheep which have been in contact with infected sheep are treated.

Contact between treated infected and non-treated, non-infected flocks must be avoided until at least 7 days after the last treatment.

Special precautions to be taken by the person administering the veterinary medicinal product to the animals:

Do not smoke, eat or drink while handling the veterinary medicinal product. Wash hands after use.

Take care to avoid self-injection as the veterinary medicinal product may cause local irritation and/or pain at the injection site. In case of accidental self-injection, seek medical advice and show the label or package leaflet to the physician.

Special precautions for the protection of the environment:

Not applicable

3.6 Adverse events

Cattle:

Rare	Injection site swelling (soft tissue) ¹
(1 to 10 animals / 10,000 animals treated):	
Very rare	Discomfort ²
(<1 animal / 10,000 animals treated, including isolated reports):	

¹ disappears without treatment

Sheep:

Very rare:	Pain ¹
(<1 animal / 10,000 animals treated, including isolated reports):	

¹ sometimes intense but usually transient; disappears without treatment

Pigs:

Very rare:	Pain ¹
(<1 animal / 10,000 animals treated, including isolated reports):	

¹ mild and transient; but disappears without treatment

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a veterinarian, to either the marketing authorisation holder or the national competent authority via the national reporting system. See the package leaflet for respective contact details.

3.7 Use during pregnancy, lactation or lay

The veterinary medicinal product can be administered to beef cows and ewes at any stage of pregnancy or lactation provided that the milk is not intended for human consumption, and to sows at any stage of pregnancy or lactation.

The veterinary medicinal product will not affect the fertility of breeding ewes, rams, sows and boars. The veterinary medicinal product can be given to all ages of animals including young calves, lambs and piglets.

² transient after subcutaneous administration

3.8 Interaction with other medicinal products and other forms of interaction

The veterinary medicinal product has been used concurrently without adverse effects with foot and mouth disease vaccine or clostridial vaccine, given at separate injection sites.

Adequate vaccination of sheep against clostridial infections is strongly recommended.

3.9 Administration routes and dosage

The veterinary medicinal product should be given only by subcutaneous injection at the recommended dosage level of 200 mcg ivermectin per kg bodyweight under the loose skin in front of, or behind, the shoulder in cattle and in the neck in sheep. At the recommended dosage level of 300 mcg ivermectin per kg of bodyweight, the veterinary medicinal product should be given only subcutaneously in the neck in pigs.

Syringes must be filled from the vial through a dry sterile draw-off needle that has been placed in the vial stopper. Vial stoppers must not be broached more than 20 times.

This product does not contain an antimicrobial preservative. Swab septum before removing each dose.

Use sterile needle and syringe. When treating groups of animals use only an automatic dosing device (with vented draw-off apparatus when using the 50ml vial).

To ensure administration of a correct dose, body weight should be determined as accurately as possible; accuracy of the dosing device should be checked.

Use this chart as a guide in working out the appropriate dose rate:

Cattle (1 ml/50 kg)		Sheep (0.5 ml/25 kg)		, •	
Bodyweight	Dose	Bodyweight	Dose	Bodyweight	Dose
(kg)	Volume	(kg)	Volume	(kg)	Volume (ml)
	(ml)		(ml)		
Up to 50	1.0	Up to 5	0.1	Less than 4	0.1
51 – 100	2.0	5.1 – 10	0.2	5 – 7	0.2
101 – 150	3.0	10.1 – 15	0.3	8 – 10	0.3
151 – 200	4.0	15.1 – 25	0.5	11 – 13	0.4
201 – 250	5.0	25.1 – 50	1.0	14 – 16	0.5
251 – 300	6.0	50.1 – 75	1.5	17 – 33	1.0
301 – 350	7.0	75.1 - 100	2.0	34 – 50	1.5
351 - 400	8.0			51 – 66	2.0
				67 – 99	3.0
				100 – 133	4.0
				134 – 166	5.0
				167 - 200	6.0
For cattle weighing		For sheep we	eighing	For pigs weig	ghing over
over 400 kg calculate		over 100 kg calculate		200 kg calcu	late the dose
the dose at the rate of		the dose at the rate		at the rate of	1 ml per 33
1 ml per 50 kg		of 0.5 ml per	25 kg	kg bodyweig	ht.
bodyweight.		bodyweight.			

	l I

When treating pigs and sheep of less than 16 kg, seek veterinary advice regarding the use of 1ml disposable syringes graduated in increments of 0.1 ml. When treating individual sheep, a syringe, not exceeding 2 ml and calibrated in increments of 0.1 ml, should be used.

Each ml contains 10 mg of ivermectin sufficient to treat 50 kg of bodyweight of cattle and sheep and 33 kg of bodyweight of pigs. The injection may be given with any standard automatic or single-dose or hypodermic syringe. Use of 17-gauge x ½ inch needle is suggested. Replace with a fresh sterile needle after every 10 to 12 animals. Injection of wet or dirty animals is not recommended.

For the treatment and control of sheep scab (*Psoroptes ovis*), two injections with a seven-day interval are required to treat clinical signs of scab and to eliminate mites.

3.10 Symptoms of overdose (and where applicable, emergency procedures and antidotes)

Cattle

Single doses of 4 mg ivermectin per kg (20x the recommended use level) given subcutaneously resulted in ataxia and depression.

Sheep

At oral dose levels up to 4 mg ivermectin per kg (20x the recommended use level) given subcutaneously resulted in ataxia and depression.

Pigs

A dose of 30 mg ivermectin per kg (100x the recommended dose of 0.3 mg per kg) injected subcutaneously to pigs caused lethargy, ataxia, bilateral mydriasis, intermittent tremors, laboured breathing and lateral recumbency.

No antidote has been identified; however, symptomatic therapy may be beneficial.

3.11 Special restrictions for use and special conditions for use, including restrictions on the use of antimicrobial and antiparasitic veterinary medicinal products in order to limit the risk of development of resistance

Not applicable.

3.12 Withdrawal periods

Cattle (meat & offal): 49 days.

Cattle (milk): Do not use in lactating cows producing milk for human consumption. Do not use in non-lactating dairy cows, including pregnant heifers, within 60 days of calving.

Sheep (meat & offal): 37 days.

Sheep (milk): Do not use in lactating sheep producing milk for human consumption.

Pigs (meat & offal): 19 days.

4. PHARMACOLOGICAL INFORMATION

4.1 ATC Vet Code: QP54AA01

4.2 Pharmacodynamics

Mechanism of Action

Ivermectin is a member of the macrocyclic lactone class of endectocides which have a unique mode of action. Compounds of the class bind selectively and with high affinity to glutamate-gated chloride ion channels which occur in invertebrate nerve and muscle cells. This leads to an increase in the permeability of the cell membrane to chloride ions with hyperpolarization of the nerve or muscle cell, resulting in paralysis and death of the parasite. Compounds of this class may also interact with other ligand-gated chloride channels, such as those gated by the neurotransmitter gamma-aminobutyric acid (GABA).

The margin of safety for compounds of this class is attributable to the fact that mammals do not have glutamate-gated chloride channels, the macrocyclic lactones have a low affinity for other mammalian ligand-gated chloride channels and they do not readily cross the blood-brain barrier.

4.3 Pharmacokinetics

Maximum plasma concentration Cattle:

At a dose level of 0.2 mg ivermectin per kg a maximum plasma concentration of 35-50 ng/ml is reached in +/- 2 days and the half-life in plasma is 2.8 days. It is also established that ivermectin is carried mainly in the plasma (80 %). This distribution between plasma and blood cells remain relatively constant.

Sheen:

At a dose level of 0.3 mg ivermectin per kg an average peak of 16 ng/ml is reached one day after injection.

Pigs

During trials carried out at a dose rate of 0.2 mg/kg ivermectin, a

plasma concentration of 10-20 ng/ml was reached in +/- 2 days and half-life in plasma was 0.5 day.

Excretion: length of time and route Cattle:

A liquid chromatographic method with fluorescence detection allows the determination of ivermectin residues in tissues. After an injection of 0.3 mg ivermectin per kg, the liver (target tissue) had residues ranging from 454 ppb at 2 days post treatment to 11 ppb at 28 days post treatment.

The injection site had residues shortly after treatment, ranging up to 69 ppm at 2 days withdrawal, but by 28 days the average residue was negligible (< 2 ppb). Cattle receiving a single dose of tritium-labelled ivermectin (0.2 - 0.3 mg/kg body weight) were slaughtered at 7, 14, 21 and 28 days after dosing.

Composites of faeces collected during the first 7 days after dosing contained almost all the dosed radioactivity. Only about 1-2 % of the dosed radioactivity was excreted in the urine.

Analyses of the faeces showed that about 40-50 % of the excreted radioactivity was present as unaltered drug. The remaining 50-60 % was present as metabolites or degradation products almost all which were more polar than the ivermectin.

Sheep

A liquid chromatographic method with fluorescence detection allows the determination of ivermectin residues in tissues. After an injection of 0.3 mg ivermectin per kg, the liver (target tissue) had residues ranging from 160 ppb at 3 days post treatment to 7.2 ppb at 28 days post treatment. The highest residue levels were recovered in fat (from 230 ppb at 3 days post treatment to 13 ppb at 28 days post treatment). Residues in all tissues were below 30 ppb at 28 days post treatment. Radioactive ivermectin was administered to sheep at a dose rate of 0.3 mg per kg. Analyses of the faeces showed that about 99 % of the drug and its metabolites are excreted in the faeces, +/- 1 % being excreted in the urine.

Pigs

A liquid chromatographic method with fluorescence detection allows the determination of ivermectin residues in tissues. After an injection of 0.4 mg/kg ivermectin the liver (target tissue) contained average residues ranging from 69 ppb at 3 days post dose to 13 ppb at 14 days post dose. No liver residue (< 2 ppb) was found at 28 days post dose. Swine receiving a single dose of tritium-labelled ivermectin (0.3-0.4 mg/kg) were slaughtered at 1, 7, 14 and 28 days after dosing. Composites of faeces collected during the first 7 days after dosing contained only about 36 % of the dosed radioactivity. Less than 1 % of the dosed radioactivity was found in the urine. Analysis of the faeces showed that about 40 % of the excreted radioactivity was unaltered drug.

5. PHARMACEUTICAL PARTICULARS

5.1 Major incompatibilities

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

5.2 Shelf life

Shelf-life of the veterinary medicinal product as packaged for sale: 5 years. Shelf-life after first opening the immediate packaging: 6 months.

5.4 Special precautions for storage

Store below 30 °C.

Protect from direct sunlight.

This product does not contain any antimicrobial preservative.

Following withdrawal of the first dose, use the product within 6 months.

5.4 Nature and composition of immediate packaging

Multiple-dose rubber-capped polyethylene bottles of 50 ml, 200 ml and 500 ml containing a sterile non-aqueous solution for parenteral administration. Bottles are stoppered and then either sealed by heat or crimp-sealed with an aluminium cap.

Not all pack sizes may be marketed.

5.5 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products

Medicines should not be disposed of via wastewater.

The veterinary medical product should not enter water courses as ivermectin may be dangerous for fish and other aquatic organisms.

Do not contaminate surface waters or ditches with product or used container.

6. NAME OF THE MARKETING AUTHORISATION HOLDER

Boehringer Ingelheim Animal Health UK Limited

7. MARKETING AUTHORISATION NUMBER

Vm 08327/4193

8. DATE OF FIRST AUTHORISATION

6 April 1995

9. DATE OF THE LAST REVISION OF THE SUMMARY OF THE PRODUCT CHARACTERISTICS

September 2025

10. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCT

Veterinary medicinal product subject to prescription.

Find more product information by searching for the 'Product Information Database' on www.gov.uk.

Approved 23 October 2025

Gavin Hall