

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Inflacam 20 mg/ml solution for injection for cattle, pigs and horses

### **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

One ml contains:

Active substance

Meloxicam 20 mg

Excipient

Ethanol (96%): 159.8 mg

For the full list of excipients, see section 6.1.

### **3. PHARMACEUTICAL FORM**

Solution for injection,  
Clear yellow solution.

### **4. CLINICAL PARTICULARS**

#### **4.1 Target species**

Cattle, pigs and horses

#### **4.2 Indications for use, specifying the target species**

Cattle

For use in acute respiratory infection with appropriate antibiotic therapy to reduce clinical signs in cattle.

For use in diarrhoea in combination with oral re-hydration therapy to reduce clinical signs in calves of over one week of age and young, non-lactating cattle.

For adjunctive therapy in the treatment of acute mastitis, in combination with antibiotic therapy.

For the relief of post-operative pain following dehorning in calves.

Pigs

For use in non-infectious locomotor disorders to reduce the symptoms of lameness and inflammation.

For adjunctive therapy in the treatment of puerperal septicaemia and toxæmia (mastitis-metritis-agalactia syndrome) with appropriate antibiotic therapy.

### Horses

For use in the alleviation of inflammation and relief of pain in both acute and chronic musculo-skeletal disorders.

For the relief of pain associated with equine colic.

## **4.3 Contraindications**

See also section 4.7.

Do not use in horses less than 6 weeks of age.

Do not use in animals suffering from impaired hepatic, cardiac or renal function and haemorrhagic disorders, or where there is evidence of ulcerogenic gastrointestinal lesions.

Do not use in case of hypersensitivity to the active substance or to any of the excipients.

For the treatment of diarrhoea in cattle, do not use in animals of less than one week of age.

## **4.4 Special warnings for each target species**

Treatment of calves with Inflacam 20 minutes before dehorning reduces post-operative pain. Inflacam alone will not provide adequate pain relief during the dehorning procedure. To obtain adequate pain relief during surgery co-medication with an appropriate analgesic is needed.

## **4.5 Special precautions for use**

### Special precautions for use in animals

If adverse reactions occur, treatment should be discontinued and the advice of a veterinarian should be sought.

Avoid use in very severely dehydrated, hypovolaemic or hypotensive animals which require parenteral rehydration, as there may be a potential risk of renal toxicity.

In case of inadequate relief of pain when used in the treatment of equine colic, careful re-evaluation of the diagnosis should be made as this could indicate the need for surgical intervention.

### Special precautions to be taken by the person administering the veterinary medicinal product to animals

Accidental self-injection may give rise to pain. People with known hypersensitivity to non-steroidal anti-inflammatory drugs (NSAIDs) should avoid contact with the veterinary medicinal product. In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

## **4.6 Adverse reactions (frequency and seriousness)**

A slight transient swelling at the injection site following subcutaneous administration was observed in less than 10 % of the cattle treated in clinical studies.

In horses, a transient swelling at the injection site can occur but resolves without intervention.

In very rare cases anaphylactoid reactions, which may be serious (including fatal), may occur and should be treated symptomatically.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

#### **4.7 Use during pregnancy, lactation or lay**

##### Cattle and pigs

Can be used during pregnancy and lactation.

##### Horses

Do not use in pregnant or lactating mares.

#### **4.8 Interaction with other medicinal products and other forms of interaction**

Do not administer concurrently with glucocorticosteroids, other non-steroidal anti-inflammatory drugs or with anti-coagulant agents.

#### **4.9 Amounts to be administered and administration route**

##### Cattle

Single subcutaneous or intravenous injection at a dosage of 0.5 mg meloxicam/kg body weight (i.e. 2.5 ml/100 kg body weight) in combination with antibiotic therapy or with oral re-hydration therapy, as appropriate.

##### Pigs

Single intramuscular injection at a dosage of 0.4 mg meloxicam/kg body weight (i.e. 2.0 ml/100 kg body weight) in combination with antibiotic therapy, as appropriate. If required, a second administration of meloxicam can be given after 24 hours.

##### Horses

Single intravenous injection at a dosage of 0.6 mg meloxicam/kg body weight (i.e. 3.0 ml/100 kg body weight).

For use in the alleviation of inflammation and the relief of pain in both acute and chronic musculo – skeletal disorders, Inflacam 15 mg/ml oral suspension may be used for continuation of treatment at a dosage of 0.6 mg meloxicam/kg body weight, 24 hours after administration of the injection.

Avoid introduction of contamination during use.

#### **4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary**

In the case of overdose, symptomatic treatment should be initiated.

#### 4.11 Withdrawal periods

##### Cattle

Meat and offal: 15 days.

Milk: 5 days.

##### Pigs

Meat and offal: 5 days.

##### Horses

Meat and offal: 5 days.

Not authorised for use in horses producing milk for human consumption.

### 5. PHARMACOLOGICAL PROPERTIES

Pharmacotherapeutic group: Antiinflammatory and antirheumatic products, non-steroids (oxicams).

ATCvet code: QM01AC06.

#### 5.1 Pharmacodynamic properties

Meloxicam is a non-steroidal anti-inflammatory drug (NSAID) of the oxicam class which acts by inhibition of prostaglandin synthesis, thereby exerting anti-inflammatory, anti-exudative, analgesic and antipyretic effects. It reduces leukocyte infiltration into the inflamed tissue. To a minor extent it also inhibits collagen-induced thrombocyte aggregation. Meloxicam also has anti-endotoxic properties because it has been shown to inhibit production of thromboxane B<sub>2</sub> induced by *E. coli* endotoxin administration in calves, lactating cows and pigs.

#### 5.2 Pharmacokinetic particulars

##### Absorption

After a single subcutaneous dose of 0.5 mg meloxicam/kg, C<sub>max</sub> values of 2.1 µg/ml and 2.7 µg/ml were reached after 7.7 hours and 4 hours in young cattle and lactating cows, respectively.

After two intramuscular doses of 0.4 mg meloxicam/kg, a C<sub>max</sub> value of 1.9 µg/ml was reached after 1 hour in pigs.

##### Distribution

More than 98 % of meloxicam is bound to plasma proteins. The highest meloxicam concentrations are to be found in liver and kidney. Comparatively low concentrations are detectable in skeletal muscle and fat.

##### Metabolism

Meloxicam is predominantly found in plasma. In cattle, meloxicam is also a major excretion product in milk and bile whereas urine contains only traces of the parent compound. In pigs, bile and urine contain only traces of the parent compound. Meloxicam is metabolised to an alcohol, an acid derivative and to several polar

metabolites. All major metabolites have been shown to be pharmacologically inactive. The metabolism in horses has not been investigated.

#### Elimination

Meloxicam is eliminated with a half-life of 26 hours and 17.5 hours after subcutaneous injection in young cattle and lactating cows, respectively.

In pigs, after intramuscular administration the mean plasma elimination half-life is approximately 2.5 hours.

In horses, after intravenous injection meloxicam is eliminated with a terminal half-life of 8.5 hours.

Approximately 50 % of the administered dose is eliminated via urine and the remainder via faeces.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

- Ethanol (96%)
- Poloxamer 188
- Macrogol 400
- Glycine
- Sodium hydroxide
- Hydrochloric acid, concentrated
- Meglumine
- Water for injections.

### **6.2 Major incompatibilities**

None known.

### **6.3 Shelf life**

Shelf life of the veterinary medicinal product as packaged for sale: 5 years.

Shelf life after first opening the immediate packaging: 28 days.

### **6.4 Special precautions for storage**

This veterinary medicinal product does not require any special storage conditions.

### **6.5 Nature and composition of immediate packaging**

Cardboard box with 1 colourless glass injection vial containing 20 ml, 50 ml, 100 ml or 250 ml.

Each vial is closed with a rubber stopper and sealed with an aluminium cap.

Not all pack sizes may be marketed.

**6.6 Special precautions for the disposal of unused veterinary medicinal products or waste materials derived from the use of such products**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

**7. MARKETING AUTHORISATION HOLDER**

Chanelle Pharmaceuticals Manufacturing Ltd.  
Loughrea  
Co. Galway  
Ireland

**8. MARKETING AUTHORISATION NUMBER**


Vm 08749/5013

**9. DATE OF FIRST AUTHORISATION**

09 December 2011

**10. DATE OF REVISION OF THE TEXT**

June 2021

A handwritten signature in black ink, consisting of several loops and a long horizontal stroke at the end.

Approved: 04 June 2021