SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Hyogen emulsion for injection for pigs

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each 2 ml dose contains:

Active substance:

Inactivated

Mycoplasma hyopneumoniae 2940 strain: min. 328 ELISA Units

Adjuvants:

Light liquid paraffin 187 µl

Escherichia coli J5 LPS min. 594 - max. 38000 Endotoxin unit

Excipient:

Thiomersal 50 µg

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Emulsion for injection Off-white, homogeneous emulsion.

4. CLINICAL PARTICULARS

4.1 Target species

Pigs for fattening

4.2 Indications for use, specifying the target species

For the active immunisation of fattening pigs from 3 weeks of age to reduce the occurrence and severity of lung lesions caused by *Mycoplasma hyopneumoniae* infection.

Onset of immunity: 3 weeks Duration of immunity: 26 weeks

4.3 Contraindications

None.

4.4 Special warnings for each target species

Vaccinate healthy animals only.

The data available are not sufficient to exclude the interaction of maternally derived antibodies against *Mycoplasma hyopneumoniae* with vaccine uptake. Interaction with maternally derived antibodies is known and should be taken into consideration. It is recommended to delay vaccination in piglets with residual MDA against *Mycoplasma hyopneumoniae* at the age of 3 weeks.

4.5 Special precautions for use

Special precautions for use in animals:

Not applicable.

Special precautions to be take by the person administering the veterinary medicinal product to animals:

In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

To the user:

This veterinary medicinal product contains mineral oil. Accidental injection/self-injection may result in severe pain and swelling, particularly if injected into a joint or finger, and in rare cases could result in the loss of the affected finger if prompt medical attention is not given. If you are accidentally injected with this veterinary medicinal product, seek prompt medical advice even if only a very small amount is injected and take the package leaflet with you. If pain persists for more than 12 hours after medical examination, seek medical advice again.

To the physician:

This veterinary medicinal product contains mineral oil. Even if small amounts have been injected, accidental injection with this product can cause intense swelling, which may, for example, result in ischaemic necrosis and even the loss of a digit. Expert and PROMPT surgical attention is required and may necessitate early incision and irrigation of the injected area, especially where there is involvement of finger pulp or tendon.

Special precautions for the protection of the environment: Not applicable.

Other precautions

4.6 Adverse reactions (frequency and seriousness)

Target species: Pigs for fattening

Very common	Elevated temperature*
(>1 animal / 10 animals	
treated):	Injection site swelling**
Uncommon	Hypersensitivity reaction***
(1 to 10 animals / 1,000	
animals treated):	Vomiting***
Very rare	Anaphylactic shock****
(<1 animal / 10,000	
animals treated, including	Recumbency****
isolated reports):	-

^{*}On the day of vaccination a transient mean increase in body temperature of about 1.3oC is very common. In an individual pig this increase might reach 2oC, but in all cases body temperature is back to normal the next day.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a veterinarian, to either the marketing authorisation holder or its local representative or the national competent authority via the national reporting system. See also the last section of the package leaflet for respective contact details.

4.7 Use during pregnancy, lactation or lay

Not applicable.

4.8 Interaction with other medicinal products and other forms of interaction

Safety and efficacy data are available which demonstrate that this vaccine can be mixed with Circovac and administered to piglets at one injection site. Vaccinate piglets from 3 weeks of age.

Onset of immunity: 3 weeks after vaccination when mixed with Circovac Duration of immunity: 23 weeks when mixed with Circovac.

^{**} A local reaction at the site of injection in the form of a swelling of a diameter up to 5 cm can be very common, which can last for three days. These reactions are of transient nature and do not need further treatment.

^{***} Immediate mild hypersensitivity-like reactions may occur uncommonly after vaccination, resulting in transient clinical signs such as vomiting.

^{****} Serious anaphylactic-type reactions (shock, recumbency) which may be fatal have been reported very rarely. Such reactions require prompt symptomatic treatment.

In case of mixing with Circovac, slight and transient local reactions may occur very commonly after the administration, mainly swelling (0.5 cm - 5 cm), mild pain and redness as well as in some cases oedema. These reactions resolve spontaneously within maximum 4 days. Transient lethargy may occur very commonly on the day of vaccination which resolves spontaneously within 1-2 days. An increase in individual rectal temperature of up to 2.5°C may occur commonly lasting less than 24 hours. The above adverse reactions were observed in clinical studies.

The product literature of Circovac should be consulted before mixed administration.

No information is available on the safety and efficacy of this vaccine when used with any other veterinary medicinal product, except when mixed with Circovac. A decision to use this vaccine before or after any other veterinary medicinal product therefore needs to be made on a case by case basis.

4.9 Amount(s) to be administered and administration route

Shake well before use.

Use sterile syringe and needle, respect aseptic conditions of vaccination.

When Hyogen is used alone:

Intramuscular use.

Vaccinate pigs in the side of their neck.

Administer a single dose of 2 ml from 3 weeks of age.

When Hyogen is mixed with Circovac:

The mixed use is restricted to the 100 doses (200 ml) presentations of Hyogen and to the 100 doses presentations (50 ml of reconstituted vaccine) of Circovac.

Piglets from 3 weeks of age:

Hyogen	Circovac
100 doses (200 ml of vaccine) in 250 ml bottle	100 doses for piglets (50 ml of reconstituted
	suspension + emulsion)

Vaccine devices should be used under aseptic conditions and in accordance with the device instructions provided by the manufacturer.

Prepare Circovac by vigorously shaking the vial of antigen suspension and injecting its content into the vial of emulsion containing adjuvant.

Mix 200 ml of Hyogen and 50 ml of Circovac and shake gently until a homogeneous white emulsion is obtained.

Administer one 2.5 ml dose of the mixture by intramuscular injection, in the side of the neck.

Use the entire vaccine mixture immediately after mixing.

4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary

As the vaccine is inactivated, studies investigating the safety of an overdose administration are not required.

4.11 Withdrawal period(s)

Zero days.

5. IMMUNOLOGICAL PROPERTIES

Pharmacotherapeutic group: Pig / Inactivated bacterial vaccines /

mycoplasma

ATCvet code: QI09AB13

Inactivated bacterial vaccine, containing whole cell concentrate of *Mycoplasma hyopneumoniae* strain 2940. This antigen is incorporated in an adjuvant for stimulation of immunity, based on a combination of light liquid paraffin and cell free *Escherichia coli* J5 LPS. The product stimulates the development of active immunity in pigs against *Mycoplasma hyopneumoniae*.

Under experimental conditions reduction of *M. hyopneumoniae* colonisation was demonstrated 44-50 days post vaccination.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Light liquid paraffin Sorbitan trioleate

Polysorbate 80

Escherichia coli J5 LPS

Thiomersal

Sodium chloride

Potassium chloride

Disodium phosphate dihydrate

Potassium dihydrogen phosphate

Water for injections

6.2 Major incompatibilities

Do not mix with any other veterinary medicinal product, except Circovac

6.3 Shelf life

Shelf life of the veterinary medicinal product as packaged for sale: 2 years Shelf life after first opening the immediate packaging: 10 hours

6.4 Special precautions for storage

Store and transport refrigerated ($2\Box C - 8\Box C$). Do not freeze. Protect from light.

6.5 Nature and composition of immediate packaging

Low density polyethylene bottle of 50, 100, 200 or 250 ml volume, sealed with siliconed, oil-resistant, nitrile rubber stopper and aluminium cap, in a cardboard box.

1x50 ml (1x25 doses), 1x100 ml (1x50 doses), 1x200 ml (1x100 doses) in 200 ml bottle, 1x200 ml (1x100 doses) in 250 ml bottle, 1x250 ml (1x125 doses), 5x50 ml (5x25 doses), 5x100 ml (5x50 doses), 5x200 ml (5x100 doses) in 200 ml bottles, 5x200 ml (5x100 doses) in 250 ml bottles 5x250 ml (5x125 doses).

Not all pack sizes may be marketed.

6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products

Medicines should not be disposed of via wastewater.

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

Ceva Sante Animale 8 rue de Logrono 33500 Libourne France

8. MARKETING AUTHORISATION NUMBER

Vm 14966/5050

9. DATE OF FIRST AUTHORISATION

11 June 2015

10. DATE OF THE LAST REVISION OF THE SUMMARY OF THE PRODUCT CHARACTERISTICS

October 2025

11. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS

Veterinary medicinal product subject to prescription.

Find more product information by searching for the 'Product Information Database' or 'PID' on www.gov.uk.

Gavin Hall

Approved: 10 October 2025