

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Previon 200 mg/ml solution for injection for pigs

### **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each ml contains:

#### **Active substance:**

Iron (III) .....	200.0 mg
(as gleptoferron.....)	532.6 mg)

#### **Excipients:**

Phenol .....	5.0 mg
--------------	--------

For the full list of excipients, see section 6.1.

### **3. PHARMACEUTICAL FORM**

Solution for injection.  
Dark brown, slightly viscous solution.

### **4. CLINICAL PARTICULARS**

#### **4.1 Target species**

Pigs (piglets)

#### **4.2 Indications for use, specifying the target species**

For the prevention of iron deficiency anaemia in piglets.

#### **4.3 Contraindications**

Do not use in piglets suspected to suffer from deficiency of vitamin E and/or selenium.  
Do not use in cases of hypersensitivity to the active substance.  
Do not use in clinically diseased animals, especially not in case of diarrhoea.  
Do not administer intravenously.

#### **4.4 Special warnings for each target species**

None.

#### **4.5 Special precautions for use**

##### Special precautions for use in animals

It is advisable to stretch the skin at the injection site to minimise leakage after withdrawal of the needle. Normal aseptic injection techniques should be practised. Avoid the introduction of contamination during use.

##### Special precautions to be taken by the person administering the veterinary medicinal product to animals

People with known hypersensitivity to the active substance (gleptoferron) or with hemochromatosis should avoid contact with the veterinary medicinal product. Take care to avoid accidental self-injection and contact with mucous membranes. In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician. Wash hands after use.

#### **4.6 Adverse reactions (frequency and seriousness)**

Occasionally discolouration of the tissue and/or slight, soft swelling may be observed at the site of injection. This should disappear within a few days. Hypersensitivity reactions can occur. Deaths in piglets following the administration of parenteral iron dextran preparations associated with genetic factors or deficiency of vitamin E and/or selenium have been reported rarely.

Deaths in piglets which have been attributed to an increased susceptibility to infection due to temporary blocking of the reticuloendothelial system, have been reported very rarely.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports)

#### **4.7 Use during pregnancy, lactation or lay**

Not applicable.

#### **4.8 Interaction with other medicinal products and other forms of interaction**

The absorption of concomitantly administered oral iron may be reduced.

#### **4.9 Amounts to be administered and administration route**

Intramuscular use.

The product is administered as a single 1 ml (200 mg iron) dose by deep intramuscular injection.

Inject once between the 1st and the 3rd day of life.

The use of a multidose syringe is recommended. To refill the syringe use a draw-off needle to avoid excessive broaching of the stopper. The stopper must not be broached

more than 10 times. When treating groups of animals in one run, use a draw-off needle that has been placed in the vial stopper to avoid excess broaching of the stopper. The draw-off needle has to be removed after treatment.

#### **4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary**

Large amounts of iron administered by the parenteral route may result in transient reduced capacity of the immune system due to iron overload of lymph macrophages. Pain, inflammation reactions, abscess formation as well as persistent discolouration of muscle tissue at the injection site may occur.

Iatrogenic poisoning may result in the following signs: pale mucous membranes, haemorrhagic gastroenteritis, vomiting, tachycardia, hypotension, dyspnoea, oedema of the limbs, lameness, shock, death, liver damage. Supportive measures such as chelating agents can be used.

Iron overdose can result in gastrointestinal signs such as diarrhoea or constipation. Treat symptomatically.

#### **4.11 Withdrawal period(s)**

Meat and offal: Zero days.

### **5. PHARMACOLOGICAL PROPERTIES**

Pharmacotherapeutic group: Iron, parenteral preparations.  
ATC vet code: QB03AC

#### **5.1 Pharmacodynamic properties**

Iron is an essential micronutrient. It plays a major role in the oxygen transport of haemoglobin and myoglobin, as well as a key role in enzymes, such as cytochromes, catalases, and peroxidases.

Iron has a high recovery rate from metabolism and food ingested. Thus, deficiency occurs only very rarely in adult animals.

#### **5.2 Pharmacokinetic particulars**

After intramuscular injection, the iron complex is absorbed into the lymphatic tissue within 3 days. Here, the complex is split to release  $Fe^{3+}$  which is stored as ferritin in the main storage organs (e.g. liver, spleen and the reticuloendothelial system). In the blood, free  $Fe^{3+}$  binds to transferrin (transport form) and is mainly used for the synthesis of haemoglobin.

### **6. PHARMACEUTICAL PARTICULARS**

#### **6.1 List of excipients**

Phenol  
Water for injections

## **6.2 Major incompatibilities**

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

## **6.3 Shelf life**

Shelf life of the veterinary medicinal product as packaged for sale: 3 years.  
Shelf life after first opening the immediate packaging: 28 days.

## **6.4 Special precautions for storage**

This veterinary medicinal product does not require any special storage conditions.

## **6.5 Nature and composition of immediate packaging**

100 ml and 250 ml amber Type II glass vials, closed with Type I polymeric elastomer stopper with aluminium cap.

### Pack sizes:

Box with 1 vial of 100 ml  
Box with 1 vial of 250 ml

Not all pack sizes may be marketed.

## **6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

## **7. MARKETING AUTHORISATION HOLDER**

Laboratorios Hipra, S.A.  
Avda. la Selva, 135  
17170 Amer (Girona)  
SPAIN Tel. (34) 972 43 06 60  
Fax (34) 972 43 06 61  
E-mail: hipra@hipra.com

## **8. MARKETING AUTHORISATION NUMBER**

Vm 17533/4017

## **9. DATE OF FIRST AUTHORISATION**

09 November 2016

**10. DATE OF REVISION OF THE TEXT**

November 2021

Approved: 12/11/21

A handwritten signature in black ink that reads "D. Austin". The signature is written in a cursive style with a horizontal line extending to the right from the end of the name.