

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Sevotek 1000 mg/g inhalation vapour, liquid for dogs and cats

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each g contains:

Active substances:

Sevoflurane 1000 mg

Excipients: None

Clear, colourless liquid.

3. CLINICAL INFORMATION

3.1 Target species

Dogs and cats.

3.2 Indications for use for each target species

For the induction and maintenance of anaesthesia.

3.3 Contraindications

Do not use in cases of hypersensitivity to the active substance or other halogenated anaesthetic agents.

Do not use in animals with a known or suspected genetic susceptibility to malignant hyperthermia.

3.4 Special warnings

None.

3.5 Special precautions for use

Special precautions for safe use in the target species:

Halogenated volatile anaesthetics can react with dry carbon dioxide (CO₂) absorbents to produce carbon monoxide (CO) that may result in elevated levels of carboxyhaemoglobin in some dogs. In order to minimise this reaction in rebreathing anaesthetic circuits, the veterinary medicinal product should not be passed through soda lime or barium hydroxide that has been allowed to dry out.

The exothermic reaction that occurs between inhalation agents (including sevoflurane) and CO₂ absorbents is increased when the CO₂ absorbent becomes desiccated, such as after an extended period of dry gas flow through the CO₂ absorbent canisters. Rare cases of excessive heat production, smoke and/or fire in the anaesthetic machine have been reported during the use of a desiccated CO₂ absorbent and sevoflurane. An unusual decrease in the expected depth of anaesthesia compared to the vapouriser setting may indicate excessive heating of the CO₂ absorbent canister.

If it is suspected that the CO₂ absorbent may be desiccated, it must be replaced. The colour indicator of most CO₂ absorbents does not necessarily change as a result of desiccation. Therefore, the lack of significant colour change should not be taken as an assurance of adequate hydration. CO₂ absorbents should be replaced routinely regardless of the state of the colour indicator.

1,1,3,3,3-pentafluoro-2-(fluoromethoxy)propene (C₄H₂F₆O), also known as Compound A, is produced when sevoflurane interacts with soda lime or barium hydroxide. Reaction with barium hydroxide results in a greater production of Compound A than does the reaction with soda lime. Its concentration in a circle absorber system increases with increasing sevoflurane concentrations and with decreasing fresh gas flow rates. Sevoflurane degradation in soda lime has been shown to increase with temperature. Since the reaction of carbon dioxide with absorbents is exothermic, this temperature increase will be determined by the quantities of CO₂ absorbed, which in turn will depend on fresh gas flow in the anaesthetic circle system, metabolic status of the dog and ventilation. Although Compound A is a dose-dependent nephrotoxin in rats, the mechanism of this renal toxicity is unknown. Long-duration, low-flow sevoflurane anaesthesia should be avoided due to the risks of Compound A accumulation.

During maintenance of anaesthesia, increasing the concentration of sevoflurane produces a dose-dependent decrease in blood pressure. Due to sevoflurane's low solubility in blood, these haemodynamic changes may occur more rapidly than with other volatile anaesthetics. Arterial blood pressure should be monitored at frequent intervals during sevoflurane anaesthesia. Facilities for artificial ventilation, oxygen enrichment and circulatory resuscitation should be immediately available. Excessive decreases in blood pressure or respiratory depression may be related to the depth of anaesthesia and may be corrected by decreasing the inspired concentration of sevoflurane. The low solubility of sevoflurane also facilitates rapid elimination by the lungs. The nephrotoxic potential of certain NSAIDs, when used in the perioperative period, may be exacerbated by hypotensive episodes during sevoflurane anaesthesia. In order to maintain renal blood flow, prolonged episodes of hypotension (mean arterial pressure below 60 mmHg) should be avoided in dogs and cats during sevoflurane anaesthesia.

In common with all volatile agents, sevoflurane may cause hypotension in hypovolaemic animals such as those requiring surgery to repair traumatic injury, and lower doses should be administered in combination with appropriate analgesics.

Sevoflurane may trigger episodes of malignant hyperthermia in susceptible dogs and cats. If malignant hyperthermia develops, the anaesthetic supply should be

interrupted immediately and 100% oxygen administered using fresh anaesthetic hoses and a rebreathing bag. Appropriate treatment should readily be instituted.

Compromised or debilitated dogs and cats

Doses of sevoflurane may need adjustment for geriatric or debilitated animals. Doses required for maintenance anaesthesia may need to be reduced by approximately 0.5% in geriatric dogs (i.e. 2.8% to 3.1% in premedicated geriatric dogs and 3.2 to 3.3% in unpremedicated geriatric dogs). There is no information on the adjustment of the maintenance dose in cats. Adjustment is, therefore, left to the discretion of the veterinarian. Limited clinical experience in administering sevoflurane to animals with renal, hepatic and cardiovascular insufficiency suggests that sevoflurane may be safely used in these conditions. However, it is recommended that such animals be monitored carefully during sevoflurane anaesthesia.

Sevoflurane may cause a small increase in intracranial pressure (ICP) under conditions of normocapnia in dogs. In dogs with head injuries or other conditions placing them at risk from increased ICP, it is recommended that hypocapnia be induced by means of controlled hyperventilation as a means of preventing changes in ICP.

There are limited data to support the safety of sevoflurane in animals less than 12 weeks of age. Therefore, it should only be used in these animals according to a benefit-risk assessment by the responsible veterinary surgeon.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

In order to minimise exposure to sevoflurane vapour, the following recommendations are made:

- Use a cuffed endotracheal tube when possible for the administration of the veterinary medicinal product during maintenance anaesthesia.
- Avoid using masking procedures for prolonged induction and maintenance of general anaesthesia.
- Ensure that operating rooms and animal recovery areas are provided with adequate ventilation or scavenging systems to prevent the accumulation of anaesthetic vapour.
- All scavenging/extraction systems must be adequately maintained.
- Pregnant and breast-feeding women should not have any contact with the veterinary medicinal product and should avoid operating rooms and animal recovery areas.
- Care should be taken when dispensing the veterinary medicinal product, with immediate removal of any spillage.
- Do not inhale the vapour directly.
- Avoid contact by mouth.

- Halogenated anaesthetic agents may induce liver damage. This is an idiosyncratic response very occasionally seen after repeated exposure.
- From an environmental point of view, it is considered good practice to use charcoal filters with scavenging equipment.

Direct exposure to eyes may result in mild irritation. If eye exposure occurs, wash with plenty of water for 15 minutes. Seek medical attention if irritation persists

In case of accidental contact with the skin, wash affected area with abundant water.

Symptoms of human overexposure (inhalation) to sevoflurane vapour include respiratory depression, hypotension, bradycardia, shivering, nausea and headache. If these symptoms occur, the individual should be removed from the source of exposure and medical attention sought.

To the physician:

Maintain a patent airway and give symptomatic and supportive treatment.

Special precautions for the protection of the environment:

Not applicable.

3.6 Adverse events

Dogs, cats:

Very common (>1 animal / 10 animals treated):	Hypotension, tachypnoea, muscle tenseness, excitation, apnoea, fasciculations ¹ , emesis
Common (1 to 10 animals / 100 animals treated):	Respiratory depression ² , Bradycardia ³
Very rare (<1 animal / 10 000 animals treated, including isolated reports):	Paddling, retching, hypersalivation, cyanosis, premature ventricular contractions, cardiac depression ⁴
Undetermined frequency (cannot be estimated from the available data):	Circulatory disorder ⁵ , Malignant hyperthermia, Elevated aspartate aminotransferase (AST) ⁶ , elevated alanine aminotransferase (ALT) ⁶ , elevated lactate dehydrogenase (LDH) ^{6,7} , elevated bilirubin ^{6,7} , elevated white blood cell counts ^{6,7}

¹ Muscle

² Dose-dependent. Respiration should be closely monitored during sevoflurane anaesthesia and the inspired concentration of sevoflurane adjusted accordingly.

³ It may be reversed by administration of anticholinergics.

⁴ Excessive

⁵ Decreased renal blood flow, as a result of hypotension induced by sevoflurane anaesthesia.

⁶ Transient

⁷ Only in dogs

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a veterinarian, to either the marketing authorisation holder or the national competent authority via the national reporting system. See the package leaflet for respective contact details.

3.7 Use during pregnancy, lactation or lay

Pregnancy and lactation:

The safety of the veterinary medicinal product has not been established during pregnancy or lactation. However, there is limited clinical experience of the use of sevoflurane, after propofol induction, in bitches and queens undergoing caesarean section, without any ill effects being detected in either the bitch or queen, or the puppies or kittens. Use only according to the risk-benefit assessment of the responsible veterinarian.

3.8 Interaction with other medicinal products and other forms of interaction

Intravenous Anaesthetics:

Sevoflurane administration is compatible with the intravenous barbiturates and propofol and in cats alfaxalone and ketamine. In dogs, the concurrent administration of thiopental, however, may slightly increase sensitivity to adrenaline-induced cardiac arrhythmias.

Benzodiazepines and Opioids:

Sevoflurane administration is compatible with benzodiazepines and opioids commonly used in veterinary practice. In common with other inhalational anaesthetics, the MAC of sevoflurane is reduced by the concurrent administration of benzodiazepines and opioids.

Phenothiazines and alpha-2-agonists:

Sevoflurane is compatible with phenothiazines and alpha-2-agonists commonly used in veterinary practice. Alpha-2-agonists have an anaesthetic sparing effect and therefore the dose of sevoflurane should be reduced accordingly. Limited data are available on the effects of the highly potent alpha-2-agonists (medetomidine, romifidine and dexmedetomidine) as premedication. Therefore they should be used with caution. Alpha-2-agonists cause bradycardia which may occur when they are used with sevoflurane. Bradycardia can be reversed by the administration of anticholinergics.

Anticholinergics:

Studies in dogs and cats show that anticholinergic premedication is compatible with sevoflurane anaesthesia in dogs and cats.

In a laboratory study, the use of an acepromazine/oxymorphone/thiopental/sevoflurane anaesthetic regimen resulted in prolonged recoveries in all the dogs treated, compared to recoveries in dogs anaesthetised with sevoflurane alone.

The use of sevoflurane with nondepolarising muscle relaxants has not been evaluated in dogs. In cats sevoflurane has been shown to exert some neuromuscular blocking effect, but this is only apparent at high doses. In humans sevoflurane increases both the intensity and duration of neuromuscular blockade induced by nondepolarising muscle relaxants. Neuromuscular blocking agents have been used in cats anaesthetised with sevoflurane without any unexpected effects.

3.9 Administration routes and dosage

Inhalation use.

Inspired Concentration:

The veterinary medicinal product should be administered via a vaporiser specifically calibrated for use with sevoflurane so that the concentration delivered can be accurately controlled. The veterinary medicinal product contains no stabiliser and does not affect the calibration or operation of these vaporisers in any way. The administration of sevoflurane must be individualised based on the dog's or cat's response.

Premedication:

The necessity for and choice of premedication is left to the discretion of the veterinarian. Preanaesthetic doses for premedicants may be lower than the label directions for their use as a single medication.

Induction of anaesthesia:

For mask induction using sevoflurane, inspired concentrations of 5 to 7% sevoflurane with oxygen are employed to induce surgical anaesthesia in the healthy dog, and 6 to 8% sevoflurane with oxygen in the cat. These concentrations can be expected to produce surgical anaesthesia within 3 to 14 minutes in dogs and within 2 to 3 minutes in cats. Sevoflurane concentration for induction may be set initially, or may be achieved gradually over the course of 1 to 2 minutes. The use of premedicants does not affect the concentration of sevoflurane required for induction.

Maintenance of anaesthesia:

Sevoflurane may be used for maintenance anaesthesia following mask induction with sevoflurane or following induction with injectable agents. The concentration of sevoflurane necessary to maintain anaesthesia is less than that required for induction.

Surgical levels of anaesthesia in the healthy dog may be maintained with inhaled concentrations of 3.3 to 3.6% in the presence of premedication. In the absence of premedication, inhaled concentrations of sevoflurane in the range 3.7 to 3.8% will

provide surgical levels of anaesthesia in the healthy dog. In the cat surgical anaesthesia is maintained with sevoflurane concentrations of 3.7-4.5%. The presence of surgical stimulation may require an increase in the concentration of sevoflurane. The use of injectable induction agents without premedication has little effect on the concentrations of sevoflurane required for maintenance. Anaesthetic regimens that include opioid, alpha-2-agonist, benzodiazepine or phenothiazine premedication will allow the use of lower sevoflurane maintenance concentrations.

3.10 Symptoms of overdose (and where applicable, emergency procedures and antidotes)

Sevoflurane overdose may result in profound respiratory depression. Therefore, respiration must be monitored closely and supported when necessary with supplementary oxygen and/or assisted ventilation.

In cases of severe cardiopulmonary depression, administration of sevoflurane should be discontinued, the existence of a patent airway ensured, and assisted or controlled ventilation with pure oxygen initiated. Cardiovascular depression should be treated with plasma expanders, pressor agents, antiarrhythmic agents or other appropriate techniques.

Due to sevoflurane's low solubility in blood, increasing the concentration may result in rapid haemodynamic changes (dose-dependent decreases in blood pressure) compared to other volatile anaesthetics. Excessive decreases in blood pressure or respiratory depression may be corrected by decreasing or discontinuing the inspired concentration of sevoflurane.

3.11 Special restrictions for use and special conditions for use, including restrictions on the use of antimicrobial and antiparasitic veterinary medicinal products in order to limit the risk of development of resistance

For administration only by a veterinarian.

3.12 Withdrawal periods

Not applicable

4. PHARMACOLOGICAL INFORMATION

4.1 ATCvet code: QN01AB08

4.2 Pharmacodynamics

Sevoflurane is an inhalation anaesthetic agent, having a light odour, for induction and maintenance of general anaesthesia. The Minimum Alveolar Concentration (MAC) of sevoflurane in dogs is 2.36% and the MAC in cats 3.1%. Multiples of MAC are used as a guide for surgical levels of anaesthesia, which are typically 1.3 to 1.5 times the MAC value.

Sevoflurane produces unconsciousness by its action on the central nervous system. Sevoflurane produces only modest increases in cerebral blood flow and metabolic rate, and has little or no ability to potentiate seizures. In the dog sevoflurane may increase intracranial pressure at concentrations of 2.0 MAC and above under normal partial pressures of carbon dioxide (normocapnia), but intracranial pressure has been shown to remain within normal range at sevoflurane concentrations of up to 1.5 MAC if hypocapnia is induced by hyperventilation. Sevoflurane in the cat did not increase intracranial pressure during normocapnia.

Sevoflurane has a variable effect on heart rate, which tends to increase from baseline at low MAC and fall back with increasing MAC. Sevoflurane causes systemic vasodilation and produces dose- dependent decreases in mean arterial pressure, total peripheral resistance, cardiac output and possibly the strength of myocardial contraction and speed of myocardial relaxation.

Sevoflurane has a depressive effect on respiration characterised by a fall in ventilation frequency. Respiratory depression may lead to respiratory acidosis and respiratory arrest (at sevoflurane concentrations of 2.0 MAC and above) in spontaneously breathing dogs and cats.

In dogs concentrations of sevoflurane below 2.0 MAC result in a small net increase in total liver blood flow. Hepatic oxygen delivery and consumption were not significantly altered at concentrations up to 2.0 MAC.

Sevoflurane administration adversely affects the autoregulation of renal blood flow in dogs and cats. As a result, renal blood flow falls in a linear fashion with increasing hypotension in sevoflurane anaesthetised dogs and cats. Nevertheless, renal oxygen consumption, and hence renal function, are preserved at mean arterial pressures above 60 mmHg in dogs and cats.

In cats no effect of sevoflurane on spleen size were recorded.

4.3 Pharmacokinetics

The pharmacokinetics of sevoflurane have not been investigated in the cat. However, based on sevoflurane blood solubility comparisons, feline uptake and elimination kinetics of sevoflurane are expected to be similar to those in the dog. Clinical data for the cat indicate rapid onset of, and recovery from, sevoflurane anaesthesia.

A minimal amount of sevoflurane is required to be dissolved in the blood before the alveolar partial pressure is in equilibrium with the arterial partial pressure because of the low solubility of sevoflurane in blood (blood/gas partition coefficient at 30° C is 0.63 to 0.69). During sevoflurane induction, there is a rapid increase in alveolar concentration towards the inspired concentration, with the ratio of inspired to end-tidal concentration of sevoflurane reaching a value of 1 within 10 minutes. Anaesthetic induction is correspondingly rapid and the depth of anaesthesia changes rapidly with changes in anaesthetic concentration.

Sevoflurane is metabolised to a limited extent in the dog (1 to 5%). The principle metabolites are hexafluoroisopropanol (HFIP) with release of inorganic fluoride and CO₂. Fluoride ion concentrations are influenced by the duration of anaesthesia and the concentration of sevoflurane. Once formed, HFIP is rapidly conjugated with

glucuronic acid and eliminated as a urinary metabolite. No other metabolic pathways for sevoflurane have been identified. In dogs exposed to 4% sevoflurane for 3 hours, mean peak maximum serum fluoride concentrations of $20.0 \pm 4.8 \mu\text{mol/l}$ have been observed after 3 hours of anaesthesia. Serum fluoride fell quickly after anaesthesia ended and had returned to baseline by 24 hours post-anaesthesia.

The elimination of sevoflurane is biphasic in nature, with an initial rapid phase and a second, slower phase. Parent compound (the dominant fraction) is eliminated via the lungs. The half-life for the slow elimination phase is approximately 50 minutes. Elimination from blood is largely complete within 24 hours. The elimination time from adipose tissue is more prolonged than from the brain.

5. PHARMACEUTICAL PARTICULARS

5.1 Major incompatibilities

In absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

5.2 Shelf life

Shelf-life of the veterinary medicinal product as packaged for sale: 30 months

Shelf life after first opening the immediate packaging: 1 year

5.3 Special precautions for storage

Keep the bottle tightly closed.

5.4 Nature and composition of immediate packaging

Amber coloured glass bottle (Type III) with a polypropylene/polyethylene pilfer-proof cap and a high density polyethylene neck collar with wing ("keyed" collar), which is fitted over the cap and bottle neck.

Pack size:

Box with 1 bottle of 250 ml (containing 380 g)

5.5 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products

Medicines should not be disposed of via wastewater or household waste.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any national collection systems applicable to the veterinary medicinal product concerned.

6. NAME OF MARKETING AUTHORISATION HOLDER

Laboratorios Karizoo S.A

7. MARKETING AUTHORISATION NUMBER

Vm 31223/4007

8. DATE OF FIRST AUTHORISATION

19 February 2018

9. DATE OF THE LAST REVISION OF THE SUMMARY OF THE PRODUCT CHARACTERISTICS

March 2026

10. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCT

Veterinary medicinal product subject to prescription.

Find more product information by searching for the 'Product Information Database' on www.gov.uk

Gavin Hall
Approved: 02 April 2026