

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1. NAME OF VETERINARY MEDICINAL PRODUCT**

Vitamin B<sub>1</sub> 10% w/v Solution for Injection

### **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

**Each ml contains:**

<b>Active Substance(s)</b>	<b>%w/v</b>
----------------------------	-------------

Thiamine Hydrochloride	10.0
------------------------	------

**Excipient(s)**

Benzyl Alcohol (as preservative)	1.5
----------------------------------	-----

For a full list of excipients, see section 6.1.

### **3. PHARMACEUTICAL FORM**

Solution for injection.

A clear, colourless to greenish-yellow liquid.

### **4. CLINICAL PARTICULARS**

#### **4.1 Target Species**

Cattle and Sheep

#### **4.2 Indications for use, specifying the target species**

For the treatment of cerebrocortical necrosis in cattle and sheep and as an adjunct in metabolic disorders of cattle.

#### **4.3 Contraindications**

Do not use in cases of known hypersensitivity to the active ingredient.

#### **4.4 Special warnings for each target species**

Not applicable.

#### **4.5 Special precautions for use**

i. Special precautions for use in animals

Intravenous injections should be given slowly.  
Observe aseptic techniques

- ii. Special precautions to be taken by the person administering the veterinary product to animals.

Care should be taken to avoid accidental self-injection. In the event of accidental self-injection, consult medical advice immediately and show the doctor this label. Wash hands after use.

#### **4.6 Adverse reactions (frequency and seriousness)**

Adverse effects are not anticipated following administration of thiamine.

#### **4.7 Use during pregnancy, lactation and lay**

It is not anticipated that the use of Vitamin B<sub>1</sub> Injection will lead to any undesirable effects during Pregnancy and/or Lactation.

#### **4.8 Interaction with other medicinal products and other forms of interactions**

None Known.

#### **4.9 Amounts to be administered and administration route**

By intramuscular or slow intravenous injections

Dosage: 2.5 - 5ml per 50 kg bodyweight. Repeat every 3 hours for up to a total of 5 doses.

#### **4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary**

Thiamine is very soluble in water and excess is excreted in the urine as a pyrimidine or as unchanged material. Tolerance studies have been carried out at twice the maximum recommended dose and the product was well tolerated.

#### **4.11 Withdrawal Period(s)**

Meat-Zero days

Milk-Zero hours

### **5. PHARMACOLOGICAL PROPERTIES**

#### **Pharmacotherapeutic group:**

Vitamins, Vitamin B<sub>1</sub>, plain and in combination with vitamins B<sub>6</sub> and B<sub>12</sub>, Vitamin B<sub>1</sub>, plain.

**ATCvet code:** QA11DAO1

Vitamin B<sub>1</sub>, also known as thiamine and as aneurine, is a water soluble vitamin. Aneurine is converted in the body to aneurine pyrophosphate (cocarboxylate) which acts as a coenzyme for several decarboxylating

enzyme systems, the most important of which is decarboxylase. The enzyme is necessary for the decarboxylation of pyruvic acid, an intermediate stage in carbohydrate build-up or breakdown. When carbohydrates are a major source of energy the body requirements of aneurine increase. Deficiencies affect the nervous and gastrointestinal systems and skin.

Tissues dependent on glucose or lactate-pyruvate for energy such as the brain and heart are particularly compromised in thiamine deficiency. Thiamine deficiency may be primary, due to deficiency in the diet, or secondary, because of destruction of the vitamin in the diet by thiaminase. The principal cause of thiamine deficiency is the presence of thiamine-destroying agents which are widely distributed in nature, and produced by rumen microflora. Growing sheep are particularly susceptible, and outbreaks of cerebrocortical necrosis have been associated with sulphur in the diet.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Benzyl Alcohol  
Sodium Hydroxide  
Disodium Edetate  
Water for Injections

### **6.2 Incompatibilities**

None known.

### **6.3 Shelf Life**

Shelf-life of the veterinary medicinal product as packaged for sale: 3 years  
Shelf-life after first opening the immediate packaging: 28 days

### **6.4 Special precautions for storage**

Do not store above 25°C.  
Protect from light.  
Once broached, use vial within 28 days. Discard unused material.

### **6.5 Nature and composition of immediate packaging**

50 ml amber type II glass vial with a red bromobutyl rubber bung and plain aluminium cap.

### **6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products**

Any unused product or waste material should be disposed of in accordance with national requirements.

**7. MARKETING AUTHORISATION HOLDER**

Bimeda Animal Health Limited  
2 / 3 / 4 Airton Close  
Tallaght  
Dublin 24  
Ireland

**8. MARKETING AUTHORISATION NUMBER**

Vm 50146/4029

**9. DATE OF FIRST AUTHORISATION**

27 May 1993

**10. DATE OF REVISION OF THE TEXT**

October 2018

Approved: 25 October 2018

